

## 2014 Child Behaviors and Health in Ailinglaplap Preliminary Results

The US Department of Agriculture (USDA)-supported Children's Healthy Living Program, or CHL, has been working with your community to understand the behaviors and health of children 2-8 years of age. In January, 2014, parents, teachers, community members, and project partners gathered data in your community. A total of 20 children from the Ailinglaplap community participated in this study. Here we have some preliminary results from our joint work. With this information parents and community leaders can see where things are going well and where additional energy, leadership, and resources will need to be placed to raise the next generation of healthy, happy children.

### Child sex, age, race distribution


Among the 20 children, all had information on sex, of which 8 (40%) were girls and 12 (60%) were boys. Furthermore, 6 (30%) were of age group 2-5 years and 14 (70%) were of age group 6-8 years. All had information on race, of which 19 (95%) were Native Hawaiian/ Pacific Islander (NHPI) and 1 (5%) were more than one race. From NHPI children, 90% were Marshallese.

Potential outliers with extreme values (defined as those with a value of 3 standard deviations (sd) above or below the mean) were excluded from this report.

### Children's Body Mass Index by Category as defined by the Center for Diseases Control and Prevention (CDC)

A total of 20 children were included for this analysis. Among them, 92% were healthy weight and 5% were underweight (Figure 1).

**Figure 1. Prevalence of overweight and obesity of study children from Ailinglaplap (n=20)**



Strategies that CHL are promoting in raising healthy children include: (1) a healthy lifestyle, which encourages children to move more, sleep more and spend less time with screens; (2) a healthy diet, which encourages children to drink more water, eat more fruit and vegetables and consume fewer sugar sweetened beverages. Below we show how the children are doing on achieving some of those strategies.


### Children moderate and vigorous level of physical activity (MVPA) as measured by accelerometer

In each study community, some children were asked to wear an accelerometer for 6 days, which records the child's activity by each second. Average minutes per day of children's MVPA were obtained from the accelerometer data.

In Ailinglaplap a total of 10 children had valid accelerometer data. Among those 10 children, daily average minutes of moderate and vigorous physical activity (MVPA) were 83.8 (sd=28.7).

Eighty percent of those 10 children met the national recommendation of 60 minutes a day of MVPA (Figure 2).

**Figure 2. Percentage of children meeting national recommendation of 60 minutes a day of moderate and vigorous types of physical activity**




One in four children (20%) did not meet the national recommendation of 60 minutes a day of MVPA. General speaking, children were doing well in engaging in physical activity.

### Children’s screen time

A total of 20 children were included in the analysis of screen time. Among them, average screen time, such as watching TV, video games, or DVD, or playing active or inactive video games was 3.3 hours (sd=2.5).

While the national recommendation is for each child to spend 2 or less hours on screens every day, only 30% of our study children met this recommendation (Figure 3).

**Figure 3. Percentage of children meeting the national recommendation of each child to spend 2 or fewer hours of screen time every day**



More than two-third (70%) of children appears to be spending too much time watching screens! This is an opportunity for both parents and educators to


intervene to help children spend less time with screens.

### Children Sleep Duration

A total of 20 children were included in the analysis of sleep duration. The average number of hours of sleep per day was 10.5 (sd=1.8).

Among the six 2-5 years old, 50% met the national recommendation of sleep of between 11-13 hours daily (11-13 hours in the graph). Another 50% of children slept more than 8 hours but less than 11 hours daily (8-10 hours in the graph) (Figure 4).

**Figure 4. Percent of children 2-5 years old among 2 groups of sleep duration**



### Acknowledgements

We would like to thank all the families of participants who completed our assessments so that we could make this important information available, as well as all our community partners for working with us to improve the health of children across the Pacific.

For more information about this report or the CHL Program, contact:

Julia Alfred | 692-455-6220 | rmissasapt@gmail.com

[www.CHL-Pacific.org](http://www.CHL-Pacific.org)



United States Department of Agriculture  
National Institute of Food and Agriculture  
Agriculture and Food Research Initiative (AFRI)  
No. 2011-68001-90335